



## Measure Information

This document contains the information submitted by measure developers/stewards, but is organized according to NQF's measure evaluation criteria and process. The item numbers refer to those in the submission form but may be in a slightly different order here. In general, the item numbers also reference the related criteria (e.g., item 1b.1 relates to sub criterion 1b).

### Brief Measure Information

**NQF #: 0318**

**Corresponding Measures:**

**De.2. Measure Title:** Delivered Dose of Peritoneal Dialysis Above Minimum

**Co.1.1. Measure Steward:** Centers for Medicare & Medicaid Services

**De.3. Brief Description of Measure:** Percentage of all patient months for adult patients ( $\geq 18$  years old) whose delivered peritoneal dialysis dose was a weekly Kt/Vurea  $\geq 1.7$  (dialytic + residual).

**1b.1. Developer Rationale:** Evaluation of PD adequacy every four months for adults is critical to ensure timely dose adjustment as needed, and adequate dialysis doses (Kt/Vurea  $> 1.7$  for adult patients and Kt/Vurea  $> 1.8$  for pediatric patients) have been linked to improved patient outcomes. Therefore, continued implementation of this measure is needed to ensure frequent adequacy measurement and adequate dialysis dosing.

Studies have shown a Kt/V of 1.8/week or greater in adult PD patients was associated with better serum albumin levels[1] and improved survival [2]. The ADEMEX did not show clinical benefit with in weekly Kt/V doses exceeding 1.7/week in adult CAPD patients [1].

1.Paniagua R, Amato D, Vonesh E, et al. "Effects of increased peritoneal clearances on mortality rates in peritoneal dialysis: ADEMEX, a prospective, randomized, controlled trial." Journal of the American Society of Nephrology: JASN (2002) 13:1307-20. PMID: 11961019.

2.Lo WK, Lui SL, Chan TM, et al. "Minimal and optimal peritoneal Kt/V targets: Results of an anuric peritoneal dialysis patient's survival analysis." Kidney international (2005) 67:2032-8. PMID: 15840054.

**S.4. Numerator Statement:** Number of patient months in the denominator whose delivered peritoneal dialysis was a weekly Kt/Vurea  $\geq 1.7$  (dialytic + residual, measured in the last 4 months).

**S.6. Denominator Statement:** To be included in the denominator for a particular reporting month, the patient must be on peritoneal dialysis for the entire month, be  $\geq 18$  years old at the beginning of the month, must have had ESRD for greater than 90 days at the beginning of the month, and must be assigned to that facility for the entire month.

**S.8. Denominator Exclusions:** Exclusions that are implicit in the denominator definition include

- 1) Patients not on peritoneal dialysis for the entire month
- 2) Pediatric patients ( $< 18$  years old)
- 3) Patients who have had ESRD for  $< 91$  days
- 4) Patients not assigned to the facility for the entire month

There are no additional exclusions for this measure.

**De.1. Measure Type:** Outcome: Intermediate Clinical Outcome

**S.17. Data Source:** Claims, Registry Data

**S.20. Level of Analysis:** Facility

**IF Endorsement Maintenance – Original Endorsement Date:** Nov 15, 2007 **Most Recent Endorsement Date:** Oct 02, 2015

**IF this measure is included in a composite, NQF Composite#/title:**

**IF this measure is paired/grouped, NQF#/title:**

**De.4. IF PAIRED/GROUPED, what is the reason this measure must be reported with other measures to appropriately interpret**

results? N/A

## 1. Evidence, Performance Gap, Priority – Importance to Measure and Report

Extent to which the specific measure focus is evidence-based, important to making significant gains in healthcare quality, and improving health outcomes for a specific high-priority (high-impact) aspect of healthcare where there is variation in or overall less-than-optimal performance. **Measures must be judged to meet all sub criteria to pass this criterion and be evaluated against the remaining criteria.**

### 1a. Evidence to Support the Measure Focus – See attached Evidence Submission Form

[0318\\_Evidence.docx](#)

#### 1a.1 For Maintenance of Endorsement: Is there new evidence about the measure since the last update/submission?

Do not remove any existing information. If there have been any changes to evidence, the Committee will consider the new evidence. Please use the most current version of the evidence attachment (v7.1). Please use red font to indicate updated evidence.

No

### 1b. Performance Gap

Demonstration of quality problems and opportunity for improvement, i.e., data demonstrating:

- considerable variation, or overall less-than-optimal performance, in the quality of care across providers; and/or
- Disparities in care across population groups.

**1b.1. Briefly explain the rationale for this measure** (e.g., how the measure will improve the quality of care, the benefits or improvements in quality envisioned by use of this measure)

*If a COMPOSITE (e.g., combination of component measure scores, all-or-none, any-or-none), SKIP this question and answer the composite questions.*

Evaluation of PD adequacy every four months for adults is critical to ensure timely dose adjustment as needed, and adequate dialysis doses (Kt/Vurea > 1.7 for adult patients and Kt/Vurea > 1.8 for pediatric patients) have been linked to improved patient outcomes. Therefore, continued implementation of this measure is needed to ensure frequent adequacy measurement and adequate dialysis dosing.

Studies have shown a Kt/V of 1.8/week or greater in adult PD patients was associated with better serum albumin levels[1] and improved survival [2]. The ADEMEX did not show clinical benefit with in weekly Kt/V doses exceeding 1.7/week in adult CAPD patients [1].

1.Paniagua R, Amato D, Vonesh E, et al. "Effects of increased peritoneal clearances on mortality rates in peritoneal dialysis: ADEMEX, a prospective, randomized, controlled trial." Journal of the American Society of Nephrology: JASN (2002) 13:1307-20. PMID: 11961019.

2.Lo WK, Lui SL, Chan TM, et al. "Minimal and optimal peritoneal Kt/V targets: Results of an anuric peritoneal dialysis patient's survival analysis." Kidney international (2005) 67:2032-8. PMID: 15840054.

**1b.2. Provide performance scores on the measure as specified (current and over time) at the specified level of analysis.** *(This is required for maintenance of endorsement. Include mean, std dev, min, max, interquartile range, scores by decile. Describe the data source including number of measured entities; number of patients; dates of data; if a sample, characteristics of the entities include.) This information also will be used to address the sub-criterion on improvement (4b1) under Usability and Use.*

Analysis of CROWNWeb and Medicare claims data from January to December 2017 indicated the mean percentage of patients with PD adequacy measurements that achieved the target at least once in four months was 90.8% (SD=10.9%). Distribution: Min=0%, Max=100%, 25th percentile = 71.2%, 50th percentile =94.1%, 75th percentile = 97.2%. These results indicate that on average, facilities are meeting the Kt/Vurea guidelines in 91% of PD patients. A description of the data is included in questions 1.1-1.7 under "Scientific Acceptability".

**1b.3. If no or limited performance data on the measure as specified is reported in 1b2, then provide a summary of data from the literature that indicates opportunity for improvement or overall less than optimal performance on the specific focus of measurement.**

N/A

**1b.4. Provide disparities data from the measure as specified (current and over time) by population group, e.g., by race/ethnicity,**

**gender, age, insurance status, socioeconomic status, and/or disability.** *(This is required for maintenance of endorsement. Describe the data source including number of measured entities; number of patients; dates of data; if a sample, characteristics of the entities included.) For measures that show high levels of performance, i.e., “topped out”, disparities data may demonstrate an opportunity for improvement/gap in care for certain sub-populations. This information also will be used to address the sub-criterion on improvement (4b1) under Usability and Use.*

Disparity analyses were performed among the eligible PD patients in facilities with 11 or more PD patients (n=57,969 ) to examine the difference in performance scores by sex, race, ethnicity, age, insurance status and nursing home status.

In particular, for each facility, the percent of patient-months by demographic group (sex, race, ethnicity, age, insurance status and nursing home status) was calculated. Then, the facilities were divided into quintiles (Q1-Q5) based on the percentage of patient-months in the particular demographic category (i.e., a facility with percentage of females similar to the national median will be included in quintile 3). The top 20% of facilities in terms of rank, based on the percentages of females, were classified as Q5, while the bottom 20% of facilities were classified as Q1. Average (mean) performance for the measure was calculated for each quintile, and the means were examined for trend across quintiles (Q1-Q5). A test for trend was performed to assess disparities in performance scores. There were no increasing (or decreasing) linear trends for each group across quintiles. All the test results imply that we do not have enough evidence to prove that the performance scores will increase (or decrease) as the respective percentage of demographic group increase.

The mean performance scores for percent of patient-months with a PD Kt/v measurement in each quintile, by demographic group, are presented below. Males, non-Black, non-White, non-Hispanic, Age 18-64, non-dual eligible, and non-nursing home are the respective reference categories.

Facility Level Quintiles by Population Group (Quintile 1-5):

Females (Q1=90.4%, Q2=90.9%, Q3=91.1%, Q4=90.2%, Q5=91.5%)  
Black (Q1=90.9%, Q2=91.8%, Q3= 91.2%, Q4=90.0% %, Q5=90.3%)  
White (Q1=89.8%, Q2=90.1%, Q3=90.4%, Q4=92.1%, Q5=91.8%)  
Hispanic (Q1=91.5%, Q2=92.2%, Q3=91.6%, Q4=89.9%, Q5=89.3%)  
Age 65+ (Q1=89.6%, Q2=90.6%, Q3=91.3%, Q4=91.2%, Q5=91.5%)  
Dual Eligible (Q1=90.7%, Q2=90.8%, Q3=91.5%, Q4=90.7%, Q5=90.5%)  
Nursing Home (Q1=90.7%, Q2=91.8%, Q3=91.1%, Q4=91.3%, Q5=89.6%)

**1b.5. If no or limited data on disparities from the measure as specified is reported in 1b.4, then provide a summary of data from the literature that addresses disparities in care on the specific focus of measurement. Include citations. Not necessary if performance data provided in 1b.4**

N/A

## 2. Reliability and Validity—Scientific Acceptability of Measure Properties

Extent to which the measure, as specified, produces consistent (reliable) and credible (valid) results about the quality of care when implemented. **Measures must be judged to meet the sub criteria for both reliability and validity to pass this criterion and be evaluated against the remaining criteria.**

**2a.1. Specifications** The measure is well defined and precisely specified so it can be implemented consistently within and across organizations and allows for comparability. eMeasures should be specified in the Health Quality Measures Format (HQMF) and the Quality Data Model (QDM).

**De.5. Subject/Topic Area** (check all the areas that apply):

Renal, Renal : End Stage Renal Disease (ESRD)

**De.6. Non-Condition Specific**(check all the areas that apply):

**De.7. Target Population Category** (Check all the populations for which the measure is specified and tested if any):

Populations at Risk

**S.1. Measure-specific Web Page** (Provide a URL link to a web page specific for this measure that contains current detailed specifications including code lists, risk model details, and supplemental materials. Do not enter a URL linking to a home page or to general information.)

N/A

**S.2a. If this is an eMeasure**, HQMF specifications must be attached. Attach the zipped output from the eMeasure authoring tool (MAT) - if the MAT was not used, contact staff. (Use the specification fields in this online form for the plain-language description of the specifications)

This is not an eMeasure Attachment:

**S.2b. Data Dictionary, Code Table, or Value Sets** (and risk model codes and coefficients when applicable) must be attached. (Excel or csv file in the suggested format preferred - if not, contact staff)

Attachment Attachment: 0318\_Code\_List-636839340387235256.xlsx

**S.2c.** Is this an instrument-based measure (i.e., data collected via instruments, surveys, tools, questionnaires, scales, etc.)? Attach copy of instrument if available.

No, this is not an instrument-based measure Attachment:

**S.2d.** Is this an instrument-based measure (i.e., data collected via instruments, surveys, tools, questionnaires, scales, etc.)? Attach copy of instrument if available.

Not an instrument-based measure

**S.3.1. For maintenance of endorsement:** Are there changes to the specifications since the last updates/submission. If yes, update the specifications for S1-2 and S4-22 and explain reasons for the changes in S3.2.

No

**S.3.2. For maintenance of endorsement,** please briefly describe any important changes to the measure specifications since last measure update and explain the reasons.

There are no changes to the measure specifications since the previous endorsement in 2015.

**S.4. Numerator Statement** (Brief, narrative description of the measure focus or what is being measured about the target population, i.e., cases from the target population with the target process, condition, event, or outcome) DO NOT include the rationale for the measure.

IF an OUTCOME MEASURE, state the outcome being measured. Calculation of the risk-adjusted outcome should be described in the calculation algorithm (S.14).

Number of patient months in the denominator whose delivered peritoneal dialysis was a weekly Kt/Vurea  $\geq 1.7$  (dialytic + residual, measured in the last 4 months).

**S.5. Numerator Details** (All information required to identify and calculate the cases from the target population with the target process, condition, event, or outcome such as definitions, time period for data collection, specific data collection items/responses, code/value sets – Note: lists of individual codes with descriptors that exceed 1 page should be provided in an Excel or csv file in required format at S.2b)

IF an OUTCOME MEASURE, describe how the observed outcome is identified/counted. Calculation of the risk-adjusted outcome should be described in the calculation algorithm (S.14).

Reporting months with weekly Kt/Vurea  $\geq 1.7$  (dialytic + residual) are counted in the numerator. If no weekly Kt/Vurea value is reported for a given patient in the reporting month, the most recent peritoneal dialysis weekly Kt/Vurea value in the prior 3 months is applied to the calculation for that month.

Missing, expired, and not performed are not counted as achieving the minimum weekly Kt/Vurea threshold.

**S.6. Denominator Statement** (Brief, narrative description of the target population being measured)

To be included in the denominator for a particular reporting month, the patient must be on peritoneal dialysis for the entire month, be  $\geq 18$  years old at the beginning of the month, must have had ESRD for greater than 90 days at the beginning of the month, and must be assigned to that facility for the entire month.

**S.7. Denominator Details** (All information required to identify and calculate the target population/denominator such as definitions, time period for data collection, specific data collection items/responses, code/value sets – Note: lists of individual codes with descriptors that exceed 1 page should be provided in an Excel or csv file in required format at S.2b.)

*IF an OUTCOME MEASURE, describe how the target population is identified. Calculation of the risk-adjusted outcome should be described in the calculation algorithm (S.14).*

A treatment history file is the data source for the denominator calculation used for the analyses supporting this submission. This file provides a complete history of the status, location, and dialysis treatment modality of an ESRD patient from the date of the first ESRD service until the patient dies or the data collection cutoff date is reached. For each patient, a new record is created each time he/she changes facility or treatment modality. Each record represents a time period associated with a specific modality and dialysis facility. CROWNWeb is the primary basis for placing patients at dialysis facilities and dialysis claims are used as an additional source of information in certain situations. Information regarding first ESRD service date, death, and transplant is obtained from CROWNWeb (including the CMS Medical Evidence Form (Form CMS-2728) and the Death Notification Form (Form CMS-2746)) and Medicare claims, as well as the Organ Procurement and Transplant Network (OPTN).

**S.8. Denominator Exclusions** (Brief narrative description of exclusions from the target population)

Exclusions that are implicit in the denominator definition include

- 1) Patients not on peritoneal dialysis for the entire month
- 2) Pediatric patients (<18 years old)
- 3) Patients who have had ESRD for <91 days
- 4) Patients not assigned to the facility for the entire month

There are no additional exclusions for this measure.

**S.9. Denominator Exclusion Details** (All information required to identify and calculate exclusions from the denominator such as definitions, time period for data collection, specific data collection items/responses, code/value sets – Note: lists of individual codes with descriptors that exceed 1 page should be provided in an Excel or csv file in required format at S.2b.)

N/A

**S.10. Stratification Information** (Provide all information required to stratify the measure results, if necessary, including the stratification variables, definitions, specific data collection items/responses, code/value sets, and the risk-model covariates and coefficients for the clinically-adjusted version of the measure when appropriate – Note: lists of individual codes with descriptors that exceed 1 page should be provided in an Excel or csv file in required format with at S.2b.)

N/A

**S.11. Risk Adjustment Type** (Select type. Provide specifications for risk stratification in measure testing attachment)

No risk adjustment or risk stratification

If other:

**S.12. Type of score:**

Rate/proportion

If other:

**S.13. Interpretation of Score** (Classifies interpretation of score according to whether better quality is associated with a higher score, a lower score, a score falling within a defined interval, or a passing score)

Better quality = Higher score

**S.14. Calculation Algorithm/Measure Logic** (Diagram or describe the calculation of the measure score as an ordered sequence of steps including identifying the target population; exclusions; cases meeting the target process, condition, event, or outcome; time period for data, aggregating data; risk adjustment; etc.)

Denominator: For the reporting month, patients are included in the denominator if:

Patient modality is indicated as PD during the entire month

Patient age as of the beginning of the reporting month is at least 18 years

Patient has had ESRD for greater than 90 days at the beginning of the month

Patient has been assigned to the facility for the entire month

Numerator: For the reporting month, patients from the denominator are also included in the numerator if they have a weekly Kt/Vurea  $\geq 1.7$ .

If no weekly Kt/Vurea value is reported for a given patient in a month, the most recent peritoneal dialysis weekly Kt/Vurea value in the prior 3 months is applied to the calculation for that month.

**S.15. Sampling** (If measure is based on a sample, provide instructions for obtaining the sample and guidance on minimum sample size.)

IF an instrument-based performance measure (e.g., PRO-PM), identify whether (and how) proxy responses are allowed.

N/A

**S.16. Survey/Patient-reported data** (If measure is based on a survey or instrument, provide instructions for data collection and guidance on minimum response rate.)

Specify calculation of response rates to be reported with performance measure results.

N/A

**S.17. Data Source** (Check ONLY the sources for which the measure is SPECIFIED AND TESTED).

If other, please describe in S.18.

Claims, Registry Data

**S.18. Data Source or Collection Instrument** (Identify the specific data source/data collection instrument (e.g. name of database, clinical registry, collection instrument, etc., and describe how data are collected.)

IF instrument-based, identify the specific instrument(s) and standard methods, modes, and languages of administration.

For the analyses supporting this submission, the measure is calculated using CROWNWeb as the primary data source for the Kt/V values used to determine the numerator. If a patient's Kt/V data are missing in CROWNWeb, Kt/V values from Medicare claims are used as an additional source for obtaining that information. Please see the attached data dictionary for a list of specific data elements that are used from each data source.

**S.19. Data Source or Collection Instrument** (available at measure-specific Web page URL identified in S.1 OR in attached appendix at A.1)

No data collection instrument provided

**S.20. Level of Analysis** (Check ONLY the levels of analysis for which the measure is SPECIFIED AND TESTED)

Facility

**S.21. Care Setting** (Check ONLY the settings for which the measure is SPECIFIED AND TESTED)

Other

If other: Dialysis Facility

**S.22. COMPOSITE Performance Measure** - Additional Specifications (Use this section as needed for aggregation and weighting rules, or calculation of individual performance measures if not individually endorsed.)

N/A

## 2. Validity – See attached Measure Testing Submission Form

0318\_testing\_01072019.docx

### 2.1 For maintenance of endorsement

Reliability testing: If testing of reliability of the measure score was not presented in prior submission(s), has reliability testing of the measure score been conducted? If yes, please provide results in the Testing attachment. Please use the most current version of the testing attachment (v7.1). Include information on all testing conducted (prior testing as well as any new testing); use red font to indicate updated testing.

Yes

### 2.2 For maintenance of endorsement

Has additional empirical validity testing of the measure score been conducted? If yes, please provide results in the Testing attachment. Please use the most current version of the testing attachment (v7.1). Include information on all testing conducted (prior

testing as well as any new testing); use red font to indicate updated testing.

Yes

### 2.3 For maintenance of endorsement

*Risk adjustment: For outcome, resource use, cost, and some process measures, risk-adjustment that includes social risk factors is not prohibited at present. Please update sections 1.8, 2a2, 2b1,2b4.3 and 2b5 in the Testing attachment and S.140 and S.11 in the online submission form. NOTE: These sections must be updated even if social risk factors are not included in the risk-adjustment strategy. You MUST use the most current version of the Testing Attachment (v7.1) -- older versions of the form will not have all required questions.*

No - This measure is not risk-adjusted

## 3. Feasibility

Extent to which the specifications including measure logic, require data that are readily available or could be captured without undue burden and can be implemented for performance measurement.

### 3a. Byproduct of Care Processes

For clinical measures, the required data elements are routinely generated and used during care delivery (e.g., blood pressure, lab test, diagnosis, medication order).

#### 3a.1. Data Elements Generated as Byproduct of Care Processes.

generated by and used by healthcare personnel during the provision of care, e.g., blood pressure, lab value, medical condition  
If other:

### 3b. Electronic Sources

The required data elements are available in electronic health records or other electronic sources. If the required data are not in electronic health records or existing electronic sources, a credible, near-term path to electronic collection is specified.

**3b.1. To what extent are the specified data elements available electronically in defined fields (i.e., data elements that are needed to compute the performance measure score are in defined, computer-readable fields)** Update this field for maintenance of endorsement.

ALL data elements are in defined fields in a combination of electronic sources

**3b.2. If ALL the data elements needed to compute the performance measure score are not from electronic sources, specify a credible, near-term path to electronic capture, OR provide a rationale for using other than electronic sources.** For maintenance of endorsement, if this measure is not an eMeasure (eCQM), please describe any efforts to develop an eMeasure (eCQM).

N/A

**3b.3. If this is an eMeasure, provide a summary of the feasibility assessment in an attached file or make available at a measure-specific URL. Please also complete and attach the NQF Feasibility Score Card.**

Attachment:

### 3c. Data Collection Strategy

Demonstration that the data collection strategy (e.g., source, timing, frequency, sampling, patient confidentiality, costs associated with fees/licensing of proprietary measures) can be implemented (e.g., already in operational use, or testing demonstrates that it is ready to put into operational use). For eMeasures, a feasibility assessment addresses the data elements and measure logic and demonstrates the eMeasure can be implemented or feasibility concerns can be adequately addressed.

**3c.1. Required for maintenance of endorsement. Describe difficulties (as a result of testing and/or operational use of the measure) regarding data collection, availability of data, missing data, timing and frequency of data collection, sampling, patient confidentiality, time and cost of data collection, other feasibility/implementation issues.**

**IF instrument-based, consider implications for both individuals providing data (patients, service recipients, respondents) and those whose performance is being measured.**

Data collection is accomplished via CROWNWeb, a web-based and electronic batch submission platform maintained and operated by CMS contractors. Measures reported on DFC are reviewed on a regular basis by dialysis facility providers and rare instances of

inaccurate or missing data are present based on comments reported in the DFC ticketing system.

**3c.2. Describe any fees, licensing, or other requirements to use any aspect of the measure as specified (e.g., value/code set, risk model, programming code, algorithm).**

N/A

## 4. Usability and Use

Extent to which potential audiences (e.g., consumers, purchasers, providers, policy makers) are using or could use performance results for both accountability and performance improvement to achieve the goal of high-quality, efficient healthcare for individuals or populations.

### 4a. Accountability and Transparency

Performance results are used in at least one accountability application within three years after initial endorsement and are publicly reported within six years after initial endorsement (or the data on performance results are available). If not in use at the time of initial endorsement, then a credible plan for implementation within the specified timeframes is provided.

#### 4.1. Current and Planned Use

*NQF-endorsed measures are expected to be used in at least one accountability application within 3 years and publicly reported within 6 years of initial endorsement in addition to performance improvement.*

Specific Plan for Use	Current Use (for current use provide URL)
	Public Reporting Dialysis Facility Compare <a href="http://www.medicare.gov/dialysisfacilitycompare/">http://www.medicare.gov/dialysisfacilitycompare/</a> Dialysis Facility Compare <a href="http://www.medicare.gov/dialysisfacilitycompare/">http://www.medicare.gov/dialysisfacilitycompare/</a>
	Payment Program ESRD QIP <a href="http://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/ESRDQIP/">http://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/ESRDQIP/</a> ESRD QIP <a href="http://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/ESRDQIP/">http://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/ESRDQIP/</a>

#### 4a1.1 For each CURRENT use, checked above (update for maintenance of endorsement), provide:

- Name of program and sponsor
- Purpose
- Geographic area and number and percentage of accountable entities and patients included
- Level of measurement and setting

#### DFC:

**Purpose:** Dialysis Facility Compare helps patients find detailed information about Medicare-certified dialysis facilities. They can compare the services and the quality of care that facilities provide.

**Geographic area:** United States

**Number of accountable entities:** All Medicare-certified dialysis facilities who are eligible for the measure, and have at least 11 patients (due to public reporting requirements). For the most recent update to Dialysis Facility Compare (January 2019), 1877 facilities had a score reported.

**Patients included:** All patients who meet the requirements to be included in the measure from included facilities.

#### QIP:

**Purpose:** The ESRD QIP will reduce payments to ESRD facilities that do not meet or exceed certain performance standards. The measure was added to the program for PY2015. In PY2019, the QIP began reporting a comprehensive Kt/V measure, for which this

the data used in this measure is counted. For the purposes of this review, we are considering this an active implementation of this measure.

Geographic area: United States

Number of accountable entities: All Medicare-certified dialysis facilities who are eligible for the measure, and have at least 11 patients (due to public reporting requirements). For the most recent QIP report (PY 2019), this was 6835 facilities. Since the QIP reports a comprehensive Kt/V measure, the number of facilities counted here is larger than for DFC.

Patients included: All patients who meet the requirements to be included in the measure from included facilities.

**4a1.2. If not currently publicly reported OR used in at least one other accountability application (e.g., payment program, certification, licensing) what are the reasons? (e.g., Do policies or actions of the developer/steward or accountable entities restrict access to performance results or impede implementation?)**

N/A

**4a1.3. If not currently publicly reported OR used in at least one other accountability application, provide a credible plan for implementation within the expected timeframes -- any accountability application within 3 years and publicly reported within 6 years of initial endorsement. (Credible plan includes the specific program, purpose, intended audience, and timeline for implementing the measure within the specified timeframes. A plan for accountability applications addresses mechanisms for data aggregation and reporting.)**

N/A

**4a2.1.1. Describe how performance results, data, and assistance with interpretation have been provided to those being measured or other users during development or implementation.**

**How many and which types of measured entities and/or others were included? If only a sample of measured entities were included, describe the full population and how the sample was selected.**

Results of this measure are currently reported on Dialysis Facility Compare and in the ESRD Quality Incentive Program (via the comprehensive Kt/V measure described above). All Medicare-certified dialysis facilities are eligible for reporting in both programs (approximately 7,000 dialysis facilities). Each program has a helpdesk and supporting documentation available to assist with interpretation of the measure results.

The measure developer (UM-KECC) produces and distributes the DFC data under contract with CMS. Other CMS contractors calculate and distribute the ESRD QIP measure results.

**4a2.1.2. Describe the process(es) involved, including when/how often results were provided, what data were provided, what educational/explanatory efforts were made, etc.**

For DFC, the results are first reported to facilities via a closed preview period, where facilities can review their data prior to each of the quarterly updates of the public facing Dialysis Facility Compare website. These preview reports are posted on dialysisdata.org, where facilities can also find a detailed Guide to the Quarterly Dialysis Facility Compare Reports and other supporting documentation. Facilities can submit comments/questions about their results at any time, and can request patient lists for their facilities during the specified preview periods.

For the ESRD QIP, results are first reported to facilities via closed preview period on an annual basis; facilities can review their data prior to the results becoming public at the end of the calendar year. These preview reports are posted on qualitynet.org, where facilities can also find supporting documentation and can submit comments/questions about their results.

A measures manual that describes the calculations for both of these programs in detail is published on the CMS website: [https://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/ESRDQIP/06\\_MeasuringQuality.html](https://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/ESRDQIP/06_MeasuringQuality.html)

**4a2.2.1. Summarize the feedback on measure performance and implementation from the measured entities and others described in 4d.1.**

**Describe how feedback was obtained.**

For DFC, feedback can be provided any time through contacting the dialysisdata.org helpdesk. Preview periods allow for specific times for facilities review and comment on measure calculations, and provide an opportunity to request a patient list.

For the ESRD QIP, feedback can be provided any time through contacting the QIP helpdesk. Preview periods allow for specific times for facilities review and comment on measure calculations. Comments can also be submitted in response to the Notice of Proposed

Rulemaking for each QIP payment year.

**4a2.2.2. Summarize the feedback obtained from those being measured.**

We reviewed the comments and questions submitted during the DFC preview periods that have taken place since the last maintenance (2016-present). Outside of questions about facility-specific results (such as questioning the Kt/V value on record for a particular patient), we receive a handful of questions each preview period regarding the measure specifications, such as the determination of dialysis modality.

Note that since UM-KECC is not the contractor responsible for the ESRD Quality Incentive Program, we do not have access to the detailed comments/requested that are submitted during the annual preview period for that program.

**4a2.2.3. Summarize the feedback obtained from other users**

We reviewed the public comments that were addressed in the ESRD QIP Final Rules (FRs) that have been published since the last endorsement (PY2019 – PY2022). Since PY 2019, the ESRD QIP has been reporting a combined Kt/V measure in order to allow for more reporting of data for pediatric and peritoneal dialysis patients. Most of the comments addressed in the rule have to do with that decision. In the FR for PY 2019, there were also a number of questions about how the comprehensive measure would be specified that were along similar lines to what is often asked via the DFC preview period.

**4a2.3. Describe how the feedback described in 4a2.2.1 has been considered when developing or revising the measure specifications or implementation, including whether the measure was modified and why or why not.**

The measure specifications have not been revised since the last maintenance cycle in 2015. Feedback received during DFC preview periods has resulted in more detailed and accurate documentation available to the public, primarily via the ESRD Measures Manual and the Guide to the Quarterly Dialysis Facility Reports.

**Improvement**

Progress toward achieving the goal of high-quality, efficient healthcare for individuals or populations is demonstrated. If not in use for performance improvement at the time of initial endorsement, then a credible rationale describes how the performance results could be used to further the goal of high-quality, efficient healthcare for individuals or populations.

**4b1. Refer to data provided in 1b but do not repeat here. Discuss any progress on improvement (trends in performance results, number and percentage of people receiving high-quality healthcare; Geographic area and number and percentage of accountable entities and patients included.)**

If no improvement was demonstrated, what are the reasons? If not in use for performance improvement at the time of initial endorsement, provide a credible rationale that describes how the performance results could be used to further the goal of high-quality, efficient healthcare for individuals or populations.

The following reports the performance scores for this measure at the yearly level for 2015 - 2017. This analysis a slight increase in performance across three years for the measure as implemented on DFC.

Year 2015:	N = 1799, Mean = 84.0%, Std Dev =13.4%, Min = 0.0%, Max = 100.0%
Year 2016:	N = 1898, Mean =89.0%, Std Dev = 11.9%, Min = 0.0%, Max = 100.0%
Year 2017:	N = 1984, Mean = 90.8%, Std Dev = 10.9%, Min = 0.0%, Max = 100.0%

**4b2. Unintended Consequences**

The benefits of the performance measure in facilitating progress toward achieving high-quality, efficient healthcare for individuals or populations outweigh evidence of unintended negative consequences to individuals or populations (if such evidence exists).

**4b2.1. Please explain any unexpected findings (positive or negative) during implementation of this measure including unintended impacts on patients.**

We have been encouraged by the magnitude of improvement in measure results after implementation noted in 4b1 above. We have not been notified of documented unintended impacts on patients as a result of measure implementation.

**4b2.2. Please explain any unexpected benefits from implementation of this measure.**

None that we are aware of, other than facility improvements over the last three reporting periods as noted in 4b1 and commented on in 4b2.1

## 5. Comparison to Related or Competing Measures

If a measure meets the above criteria and there are endorsed or new related measures (either the same measure focus or the same target population) or competing measures (both the same measure focus and the same target population), the measures are compared to address harmonization and/or selection of the best measure.

### 5. Relation to Other NQF-endorsed Measures

Are there related measures (conceptually, either same measure focus or target population) or competing measures (conceptually both the same measure focus and same target population)? If yes, list the NQF # and title of all related and/or competing measures.

Yes

#### 5.1a. List of related or competing measures (selected from NQF-endorsed measures)

0321 : Adult Kidney Disease: Peritoneal Dialysis Adequacy: Solute

#### 5.1b. If related or competing measures are not NQF endorsed please indicate measure title and steward.

### 5a. Harmonization of Related Measures

The measure specifications are harmonized with related measures;

**OR**

The differences in specifications are justified

#### 5a.1. If this measure conceptually addresses EITHER the same measure focus OR the same target population as NQF-endorsed measure(s):

Are the measure specifications harmonized to the extent possible?

No

#### 5a.2. If the measure specifications are not completely harmonized, identify the differences, rationale, and impact on interpretability and data collection burden.

In the maintenance cycle in 2011, 0318 was harmonized with 0321. Since then 0318 has been revised. The measure is not harmonized with 0321 missing values are not counted in the numerator, in order to prevent gaming of the measure.

### 5b. Competing Measures

The measure is superior to competing measures (e.g., is a more valid or efficient way to measure);

**OR**

Multiple measures are justified.

#### 5b.1. If this measure conceptually addresses both the same measure focus and the same target population as NQF-endorsed measure(s):

Describe why this measure is superior to competing measures (e.g., a more valid or efficient way to measure quality); OR provide a rationale for the additive value of endorsing an additional measure. (Provide analyses when possible.)

It is anticipated that this proposed measure will allow for assessment of a larger population given the denominator revision. Missing values are not counted in the numerator, in order to prevent gaming of the measure.

## Appendix

**A.1 Supplemental materials may be provided in an appendix.** All supplemental materials (such as data collection instrument or methodology reports) should be organized in one file with a table of contents or bookmarks. If material pertains to a specific submission form number, that should be indicated. Requested information should be provided in the submission form and required attachments. There is no guarantee that supplemental materials will be reviewed.

No appendix Attachment:

## Contact Information

Co.1 Measure Steward (Intellectual Property Owner): Centers for Medicare & Medicaid Services

Co.2 Point of Contact: Helen, Dollar-Maples, Helen.Dollar-Maples@cms.hhs.gov, 410-786-7214-

**Co.3 Measure Developer if different from Measure Steward:** [University of Michigan Kidney Epidemiology and Cost Center](#)  
**Co.4 Point of Contact:** [Casey, Parrotte](#), [parrotte@med.umich.edu](mailto:parrotte@med.umich.edu)

### Additional Information

**Ad.1 Workgroup/Expert Panel involved in measure development**  
Provide a list of sponsoring organizations and workgroup/panel members' names and organizations. Describe the members' role in measure development.

#### Measure Developer/Steward Updates and Ongoing Maintenance

**Ad.2 Year the measure was first released:** [2007](#)  
**Ad.3 Month and Year of most recent revision:** [04, 2019](#)  
**Ad.4 What is your frequency for review/update of this measure?** [Annually](#)  
**Ad.5 When is the next scheduled review/update for this measure?** [04, 2020](#)

**Ad.6 Copyright statement:** [N/A](#)  
**Ad.7 Disclaimers:** [N/A](#)

**Ad.8 Additional Information/Comments:** [After the submission of the testing attachment on January 7, we noticed a typo in 2b4.1 \(Meaningful Differences\). The description of the analysis mentions the wrong event \(hypercalcemia\), which was included in error. The description of the analysis performed is otherwise accurate.](#)

## NATIONAL QUALITY FORUM—Evidence (subcriterion 1a)

**Measure Number** (if previously endorsed): 0318

**Measure Title:** [Delivered Dose of Peritoneal Dialysis Above Minimum](#)

**IF the measure is a component in a composite performance measure, provide the title of the**

**Composite Measure here:** [Click here to enter composite measure #/ title](#)

**Date of Submission:** [4/2/2019](#)

### Instructions

- Complete 1a.1 and 1a.2 for all measures. If instrument-based measure, complete 1a.3.
- Complete **EITHER 1a.2, 1a.3 or 1a.4** as applicable for the type of measure and evidence.
- For composite performance measures:
  - A separate evidence form is required for each component measure unless several components were studied together.
  - If a component measure is submitted as an individual performance measure, attach the evidence form to the individual measure submission.
- All information needed to demonstrate meeting the evidence subcriterion (1a) must be in this form. An appendix of *supplemental* materials may be submitted, but there is no guarantee it will be reviewed.
- If you are unable to check a box, please highlight or shade the box for your response.
- Contact NQF staff regarding questions. Check for resources at [Submitting Standards webpage](#).

**Note:** The information provided in this form is intended to aid the Standing Committee and other stakeholders in understanding to what degree the evidence for this measure meets NQF's evaluation criteria.

### 1a. Evidence to Support the Measure Focus

The measure focus is evidence-based, demonstrated as follows:

- **Outcome:** <sup>3</sup> Empirical data demonstrate a relationship between the outcome and at least one healthcare structure, process, intervention, or service. If not available, wide variation in performance can be used as evidence, assuming the data are from a robust number of providers and results are not subject to systematic bias.
- **Intermediate clinical outcome:** a systematic assessment and grading of the quantity, quality, and consistency of the body of evidence <sup>4</sup> that the measured intermediate clinical outcome leads to a desired health outcome.
- **Process:** <sup>5</sup> a systematic assessment and grading of the quantity, quality, and consistency of the body of evidence <sup>4</sup> that the measured process leads to a desired health outcome.
- **Structure:** a systematic assessment and grading of the quantity, quality, and consistency of the body of evidence <sup>4</sup> that the measured structure leads to a desired health outcome.
- **Efficiency:** <sup>6</sup> evidence not required for the resource use component.
- For measures derived from [patient reports](#), evidence should demonstrate that the target population values the measured outcome, process, or structure and finds it meaningful.
- **Process measures incorporating Appropriate Use Criteria:** See NQF's guidance for evidence for measures, in general; guidance for measures specifically based on clinical practice guidelines apply as well.

### Notes

**3.** Generally, rare event outcomes do not provide adequate information for improvement or discrimination; however, serious reportable events that are compared to zero are appropriate outcomes for public reporting and quality improvement.

**4.** The preferred systems for grading the evidence are the Grading of Recommendations, Assessment, Development and Evaluation ([GRADE guidelines](#)) and/or modified GRADE.

**5.** Clinical care processes typically include multiple steps: assess → identify problem/potential problem → choose/plan intervention (with patient input) → provide intervention → evaluate impact on health status. If the measure focus is one

step in such a multistep process, the step with the strongest evidence for the link to the desired outcome should be selected as the focus of measurement. Note: A measure focused only on collecting PROM data is not a PRO-PM.

6. Measures of efficiency combine the concepts of resource use and quality (see NQF's [Measurement Framework: Evaluating Efficiency Across Episodes of Care](#); [AQA Principles of Efficiency Measures](#)).

**1a.1. This is a measure of:** (should be consistent with type of measure entered in De.1)

Outcome

Outcome: Click here to name the health outcome

Patient-reported outcome (PRO): Click here to name the PRO

*PROs include HRQoL/functional status, symptom/symptom burden, experience with care, health-related behaviors. (A PRO-based performance measure is not a survey instrument. Data may be collected using a survey instrument to construct a PRO measure.)*

Intermediate clinical outcome (e.g., lab value): [Kt/V](#)

Process: Click here to name what is being measured

Appropriate use measure: Click here to name what is being measured

Structure: Click here to name the structure

Composite: Click here to name what is being measured

**1a.2 LOGIC MODEL** Diagram or briefly describe the steps between the healthcare structures and processes (e.g., interventions, or services) and the patient's health outcome(s). The relationships in the diagram should be easily understood by general, non-technical audiences. Indicate the structure, process or outcome being measured.

The measure focus is the process of measuring peritoneal dialysis adequacy every four months for ESRD dialysis patients to assess adequate dialysis. This leads to improvement in mortality as follows: Measure PD adequacy-->Assess value-->Identify problem-->Identify treatment options-->Administer the appropriate treatment-->Impact on mortality.

**1a.3 Value and Meaningfulness:** IF this measure is derived from patient report, provide evidence that the target population values the measured **outcome, process, or structure** and finds it meaningful. (Describe how and from whom their input was obtained.)

N/A

**\*\*RESPOND TO ONLY ONE SECTION BELOW -EITHER 1a.2, 1a.3 or 1a.4) \*\***

**1a.2 FOR OUTCOME MEASURES including PATIENT REPORTED OUTCOMES - Provide empirical data demonstrating the relationship between the outcome (or PRO) to at least one healthcare structure, process, intervention, or service.**

N/A

**1a.3. SYSTEMATIC REVIEW(SR) OF THE EVIDENCE (for INTERMEDIATE OUTCOME, PROCESS, OR STRUCTURE PERFORMANCE MEASURES, INCLUDING THOSE THAT ARE INSTRUMENT-BASED) If the evidence is not based on a systematic review go to section 1a.4) If you wish to include more than one systematic review, add additional tables.**

**What is the source of the systematic review of the body of evidence that supports the performance measure? A systematic review is a scientific investigation that focuses on a specific question and uses explicit, prespecified scientific methods to identify, select, assess, and summarize the findings of similar but separate studies. It may include a quantitative synthesis (meta-analysis), depending on the available data. (IOM)**

- Clinical Practice Guideline recommendation (with evidence review)
- US Preventive Services Task Force Recommendation
- Other systematic review and grading of the body of evidence (e.g., *Cochrane Collaboration, AHRQ Evidence Practice Center*)
- Other

<p><b>Source of Systematic Review:</b></p> <ul style="list-style-type: none"> <li>• Title</li> <li>• Author</li> <li>• Date</li> <li>• Citation, including page number</li> <li>• URL</li> </ul>	<p>KDOQI Clinical Practice Guidelines and Clinical Practice Recommendations for 2006 Updates: Hemodialysis Adequacy, Peritoneal Dialysis Adequacy and Vascular Access. Am J Kidney Dis 48:S1-S322, 2006 (suppl 1).  <a href="http://www.kidney.org/professionals/KDOQI/guidelines_commentaries">http://www.kidney.org/professionals/KDOQI/guidelines_commentaries</a></p>
<p>Quote the guideline or recommendation verbatim about the process, structure or intermediate outcome being measured. If not a guideline, summarize the conclusions from the SR.</p>	<p>Adult Kt/V target:  <b>GUIDELINE 2. PERITONEAL DIALYSIS SOLUTE CLEARANCE TARGETS AND MEASUREMENTS</b></p> <p>Data from RCTs suggested that the minimally acceptable small-solute clearance for PD is less than the prior recommended level of a weekly Kt/Vurea of 2.0. Furthermore, increasing evidence indicates the importance of RKF as opposed to peritoneal small-solute clearance with respect to predicting patient survival. Therefore, prior targets have been revised as indicated next. 2.1 For patients with RKF (considered to be significant when urine volume is &gt; 100 mL/d): 2.1.1 The minimal "delivered" dose of total small-solute clearance should be a total (peritoneal and kidney) Kt/Vurea of at least 1.7 per week. (B)</p> <p>2.1.2 Total solute clearance (residual kidney and peritoneal, in terms of Kt/Vurea) should be measured within the first month after initiating dialysis therapy and at least once every 4 months thereafter. (B)</p> <p>2.1.3 If the patient has greater than 100 mL/d of residual kidney volume and residual kidney clearance is being considered as part of the patient's total weekly solute clearance goal, a 24-hour urine collection for urine volume and solute clearance determinations should be</p>

	<p>obtained at a minimum of every 2 months. (B) 2.2 For patients without RKF (considered insignificant when urine volume is =100 mL/d):</p> <p>2.2.1 The minimal "delivered" dose of total small-solute clearance should be a peritoneal Kt/Vurea of at least 1.7 per week measured within the first month after starting dialysis therapy and at least once every 4 months thereafter. (B)</p>
Grade assigned to the <b>evidence</b> associated with the recommendation with the definition of the grade	N/A
Provide all other grades and definitions from the evidence grading system	N/A
Grade assigned to the <b>recommendation</b> with definition of the grade	The guidelines for adult patients were graded B. Grade B: It is recommended that clinicians routinely follow the guideline for eligible patients. There is moderately strong evidence that the practice improves health outcomes.
Provide all other grades and definitions from the recommendation grading system	<p>The rating system defined in the KDOQI Guidelines was used to grade the strength of the Guideline recommendation. KDOQI defined grades as follows:</p> <p>Grade A: It is strongly recommended that clinicians routinely follow the guideline for eligible patients. There is strong evidence that the practice improves health outcomes.</p> <p>Grade B: It is recommended that clinicians routinely follow the guideline for eligible patients. There is moderately strong evidence that the practice improves health outcomes.</p> <p>Grade CPR: It is recommended that clinicians consider following the guideline for eligible patients. This recommendation is based on either weak evidence or on the opinions of the Work Group and reviewers that the practice might improve health outcomes.</p>
<p>Body of evidence:</p> <ul style="list-style-type: none"> <li>• Quantity – how many studies?</li> <li>• Quality – what type of studies?</li> </ul>	<p>20 studies, ranging from 1998-2004</p> <p>The KDOQI panel noted that the body of evidence shows a correlation between total solute clearance for urea and patient mortality and morbidity. Thus, this evidence supports that the delivered dose of dialysis should be measured frequently for assessment of adequate treatment, and treatment should be set accordingly. In particular, of the 20 studies considered in the body of evidence, the results from two randomized clinical trials were used to justify the KDOQI guidelines [2,3]. The results from additional observational studies also supported the KDOQI recommendations [see, e.g. 1,6].</p> <p>1. Bargman JM, Thorpe KE, Churchill DN: Relative contribution of residual renal function and peritoneal clearance to adequacy of</p>

	<p>dialysis: A reanalysis of the CANUSA Study. <i>J Am Soc Nephrol</i> 12:2158-2162, 2001</p> <ol style="list-style-type: none"> <li>2. Paniagua R, Amato D, Vonesh E, et al: Effects of increased peritoneal clearances on mortality rates in peritoneal dialysis: ADEMEX, a prospective, randomized, controlled trial. <i>J Am Soc Nephrol</i> 13:1307-1320, 2002</li> <li>3. Lo WK, Ho YW, Li CS, et al: Effect of Kt/V on survival and clinical outcome in CAPD patients in a randomized prospective study. <i>Kidney Int</i> 64:649-656, 2003</li> <li>4. Szeto CC, Wong TY, Leung CB, et al: Importance of dialysis adequacy in mortality and morbidity of Chinese CAPD patients. <i>Kidney Int</i> 58:400-407, 2000</li> <li>5. Diaz-Buxo JA, Lowrie EG, Lew NL, Zhang SM, Zhu X, Lazarus JM: Associates of mortality among peritoneal dialysis patients with special reference to peritoneal transport rates and solute clearance. <i>Am J Kidney Dis</i> 33:523-534, 1999</li> <li>6. Rocco MV, Frankenfield DL, Prowant B, Frederick P, Flanigan MJ: Risk factors for early mortality in U.S. peritoneal dialysis patients: Impact of residual renal function. <i>Perit Dial Int</i> 2002 22:371-379</li> <li>7. Termorshuizen F, Korevaar JC, Dekker FW, van Manen JG, Boeschoten EW, Krediet RT: The relative importance of residual renal function compared with peritoneal clearance for patient survival and quality of life: An analysis of the Netherlands Cooperative Study on the Adequacy of Dialysis (NECOSAD)-2. <i>Am J Kidney Dis</i> 41:1293-1302, 2003</li> <li>8. Chung SH, Heimbürger O, Stenvinkel P, Qureshi AR, Lindholm B: Association between residual renal function, inflammation and patient survival in new peritoneal dialysis patients. <i>Nephrol Dial Transplant</i> 18:590-597, 2003</li> <li>9. Jager KJ, Merkus MP, Dekker FW, et al: Mortality and technique failure in patients starting chronic peritoneal dialysis: Results of The Netherlands Cooperative Study on the Adequacy of Dialysis. NECOSAD Study group. <i>Kidney Int</i> 55:1476-1485, 1999</li> <li>10. Ates K, Nergizoglu G, Keven K, et al: Effect of fluid and sodium removal on mortality in peritoneal dialysis patients. <i>Kidney Int</i> 60:767-776, 2001</li> <li>11. Wang AY, Wang M, Woo J, et al: Inflammation, residual kidney function, and cardiac hypertrophy are interrelated and combine adversely to enhance mortality and cardiovascular death risk of peritoneal dialysis patients. <i>J Am Soc Nephrol</i> 15:2186-2194, 2004</li> </ol>
--	---

	<p>12. Szeto CC, Wong TY, Chow KM, Leung CB, Law MC, Li PK: Independent effects of renal and peritoneal clearances on the mortality of peritoneal dialysis patients. <i>Perit Dial Int</i> 24:58-64, 2004</p> <p>13. Szeto CC, Wong TY, Chow KM, et al: Impact of dialysis adequacy on the mortality and morbidity of anuric Chinese patients receiving continuous ambulatory peritoneal dialysis. <i>J Am Soc Nephrol</i> 12:355-360, 2001</p> <p>14. Bhaskaran S, Schaubel DE, Jassal SV, et al: The effect of small solute clearances on survival of anuric peritoneal dialysis patients. <i>Perit Dial Int</i> 20:181-187, 2000</p> <p>15. Rocco M, Soucie JM, Pastan S, McClellan WM: Peritoneal dialysis adequacy and risk of death. <i>Kidney Int</i> 58:446-457, 2000</p> <p>16. Lo WK, Tong KL, Li CS, et al: Relationship between adequacy of dialysis and nutritional status, and their impact on patient survival on CAPD in Hong Kong. <i>Perit Dial Int</i> 21:441-447, 2001</p> <p>17. Davies SJ, Phillips L, Russell GI: Peritoneal solute transport predicts survival on CAPD independently of residual renal function. <i>Nephrol Dial Transplant</i> 13:962-968, 1998</p> <p>18. Perez RA, Blake PG, Spanner E, et al: High creatinine excretion ratio predicts a good outcome in peritoneal dialysis patients. <i>Am J Kidney Dis</i> 36:362-367, 2000</p> <p>19. Park HC, Kang SW, Choi KH, Ha SK, Han DS, Lee HY: Clinical outcome in continuous ambulatory peritoneal dialysis patients is not influenced by high peritoneal transport status. <i>Perit Dial Int</i> 21:S80-S85, 2001 (suppl 3)</p> <p>20. Aslam N, Bernardini J, Fried L, Piraino B: Peritoneal dialysis clearance can replace residual renal function. <i>Perit Dial Int</i> 21:263- 268, 2001</p>
<p>Estimates of benefit and consistency across studies</p>	<p>In the adult population, among the studies showing any improvement in mortality in high total clearance versus low total clearance, relative risks ranged from 0.6 to 0.99. In one study, Kt/V was measured as continuous and found a relative risk of 0.94 per 0.1 mL/min increase in Kt/V (95% CI = 0.88, 1.02). The majority of the studies showed a benefit of higher total clearance in PD patients.</p>
<p>What harms were identified?</p>	<p>As described above in 1a.7.7, the majority of studies showed a benefit of higher total clearance in PD patients. Furthermore, there is little or no potential harm in assessing total urea Kt/V for PD patients.</p>
<p>Identify any new studies conducted since the SR. Do the new studies change the conclusions from the SR?</p>	<p>In May 2014, an additional literature search was performed. Additional pieces of evidence supporting the relationship between PD clearance and outcomes are included in the citations below as a result of that search.</p>

1. Krediet RT1, Struijk DG. Peritoneal changes in patients on long-term peritoneal dialysis. *Nat Rev Nephrol.* 2013 Jul;9(7):419-29. doi: 10.1038/nrneph.2013.99. Epub 2013 May 14.

Abstract: Long-term peritoneal dialysis can lead to morphological and functional changes in the peritoneum. Although the range of morphological alterations is known for the peritoneal dialysis population as a whole, these changes will not occur in every patient in the same sequence and to the same extent. Longitudinal studies are therefore required to help identify which patients might develop the changes. Although longitudinal studies using peritoneal biopsies are not possible, analyses of peritoneal effluent biomarkers that represent morphological alterations could provide insight. Longitudinal studies on peritoneal transport have been performed, but follow-up has often been too short and an insufficient number of parameters have been investigated. This Review will firstly describe peritoneal morphology and structure and will then focus on peritoneal effluent biomarkers and their changes over time. Net ultrafiltration will also be discussed together with the transport of small solutes. Data on the peritoneal transport of serum proteins show that serum protein levels do not increase to the same extent as levels of small solutes with long-term peritoneal dialysis. Early alterations in peritoneal transport must be distinguished from alterations that only develop with long-term peritoneal dialysis. Early alterations are related to vasoactive mediators, whereas later alterations are related to neoangiogenesis and fibrosis. Modern peritoneal dialysis should focus on the early detection of long-term membrane alterations by biomarkers--such as cancer antigen 125, interleukin-6 and plasminogen activator inhibitor 1--and the improved assessment of peritoneal transport.

2. Fissell R1, Schulman G, Pfister M, Zhang L, Hung AM. Novel dialysis modalities: do we need new metrics to optimize treatment? *J Clin Pharmacol.* 2012 Jan;52(1 Suppl):72S-8S. doi: 10.1177/0091270011414576.

Abstract: Delivered dose of hemodialysis has long been an important predictor of mortality. The limitations of conventional hemodialysis treatments have led to a renewed interest in more frequent and longer hemodialysis treatments. As alternative hemodialysis schedules have become more prevalent, a need for modified metrics to measure adequacy has emerged. In addition, there is an interest in finding

	<p>measures of hemodialysis adequacy that are more reliable in certain subgroups of patients, such as women, ethnic minority groups, or people with small body size. Finally, extended hemodialysis schedules suggest a need for metrics that can measure the clearance of solutes other than urea, such as middle-size molecules, and solutes for which clearance depends on intercompartmental transport across membranes. New metrics to quantify clearance in extended and alternate hemodialysis schedules are needed. As new metrics are developed, it is anticipated that they will also contribute to more accurate assessments of associations between clinical outcomes and delivered dose of dialysis in more intensive, nontraditional hemodialysis schedules. This review provides a historical prospective of dialysis dose and adequacy and describes the need for new metrics from both solute type and dialysis dose prospective as alternative hemodialysis schedules have emerged and become more prevalent.</p>
--	---

---

#### **1a.4 OTHER SOURCE OF EVIDENCE**

*If source of evidence is NOT from a clinical practice guideline, USPSTF, or systematic review, please describe the evidence on which you are basing the performance measure.*

**1a.4.1 Briefly SYNTHESIZE the evidence that supports the measure.** A list of references without a summary is not acceptable.

N/A

**1a.4.2 What process was used to identify the evidence?**

N/A

**1a.4.3. Provide the citation(s) for the evidence.**

N/A

**NATIONAL QUALITY FORUM—Measure Testing (subcriteria 2a2, 2b1-2b6)**

**Measure Number** (if previously endorsed): 0318

**Measure Title:** Delivered Dose of Peritoneal Dialysis Above Minimum

**Date of Submission:** [1/7/2019](#)

**Type of Measure:**

<input type="checkbox"/> Outcome (including PRO-PM)	<input type="checkbox"/> Composite – <b>STOP – use composite testing form</b>
<input checked="" type="checkbox"/> Intermediate Clinical Outcome	<input type="checkbox"/> Cost/resource
<input type="checkbox"/> Process (including Appropriate Use)	<input type="checkbox"/> Efficiency
<input type="checkbox"/> Structure	

**Instructions**

- Measures must be tested for all the data sources and levels of analyses that are specified. **If there is more than one set of data specifications or more than one level of analysis, contact NQF staff** about how to present all the testing information in one form.
- For all measures, sections 1, 2a2, 2b1, 2b2, and 2b4 must be completed.**
- For outcome and resource use measures, section 2b3 also must be completed.**
- If specified for **multiple data sources/sets of specifications** (e.g., claims and EHRs), section **2b5** also must be completed.
- Respond to **all** questions as instructed with answers immediately following the question. All information on testing to demonstrate meeting the subcriteria for reliability (2a2) and validity (2b1-2b6) must be in this form. An appendix for *supplemental* materials may be submitted, but there is no guarantee it will be reviewed.
- If you are unable to check a box, please highlight or shade the box for your response.
- Maximum of 25 pages (including questions/instructions; minimum font size 11 pt; do not change margins). **Contact NQF staff if more pages are needed.**
- Contact NQF staff regarding questions. Check for resources at [Submitting Standards webpage](#).
- For information on the most updated guidance on how to address social risk factors variables and testing in this form refer to the release notes for version 7.1 of the Measure Testing Attachment.

**Note:** The information provided in this form is intended to aid the Standing Committee and other stakeholders in understanding to what degree the testing results for this measure meet NQF’s evaluation criteria for testing.

**2a2. Reliability testing** <sup>10</sup> demonstrates the measure data elements are repeatable, producing the same results a high proportion of the time when assessed in the same population in the same time period and/or that the measure score is precise. For **instrument-based measures** (including PRO-PMs) **and composite performance measures**, reliability should be demonstrated for the computed performance score.

**2b1. Validity testing** <sup>11</sup> demonstrates that the measure data elements are correct and/or the measure score correctly reflects the quality of care provided, adequately identifying differences in quality. For **instrument-based measures (including PRO-PMs) and composite performance measures**, validity should be demonstrated for the computed performance score.

**2b2. Exclusions** are supported by the clinical evidence and are of sufficient frequency to warrant inclusion in the specifications of the measure; <sup>12</sup>

## AND

If patient preference (e.g., informed decisionmaking) is a basis for exclusion, there must be evidence that the exclusion impacts performance on the measure; in such cases, the measure must be specified so that the information about patient preference and the effect on the measure is transparent (e.g., numerator category computed separately, denominator exclusion category computed separately). <sup>13</sup>

**2b3. For outcome measures and other measures when indicated** (e.g., resource use):

- **an evidence-based risk-adjustment strategy** (e.g., risk models, risk stratification) is specified; is based on patient factors (including clinical and social risk factors) that influence the measured outcome and are present at start of care; <sup>14,15</sup> and has demonstrated adequate discrimination and calibration

## OR

- rationale/data support no risk adjustment/ stratification.

**2b4. Data analysis of computed measure scores demonstrates that methods for scoring and analysis of the specified measure allow for identification of statistically significant and practically/clinically meaningful <sup>16</sup> differences in performance;**

## OR

there is evidence of overall less-than-optimal performance.

**2b5. If multiple data sources/methods are specified, there is demonstration they produce comparable results.**

**2b6. Analyses identify the extent and distribution of missing data** (or nonresponse) and demonstrate that performance results are not biased due to systematic missing data (or differences between responders and nonresponders) and how the specified handling of missing data minimizes bias.

## Notes

**10.** Reliability testing applies to both the data elements and computed measure score. Examples of reliability testing for data elements include, but are not limited to: inter-rater/abstractor or intra-rater/abstractor studies; internal consistency for multi-item scales; test-retest for survey items. Reliability testing of the measure score addresses precision of measurement (e.g., signal-to-noise).

**11.** Validity testing applies to both the data elements and computed measure score. Validity testing of data elements typically analyzes agreement with another authoritative source of the same information. Examples of validity testing of the measure score include, but are not limited to: testing hypotheses that the measures scores indicate quality of care, e.g., measure scores are different for groups known to have differences in quality assessed by another valid quality measure or method; correlation of measure scores with another valid indicator of quality for the specific topic; or relationship to conceptually related measures (e.g., scores on process measures to scores on outcome measures). Face validity of the measure score as a quality indicator may be adequate if accomplished through a systematic and transparent process, by identified experts, and explicitly addresses whether performance scores resulting from the measure as specified can be used to distinguish good from poor quality. The degree of consensus and any areas of disagreement must be provided/discussed.

**12.** Examples of evidence that an exclusion distorts measure results include, but are not limited to: frequency of occurrence, variability of exclusions across providers, and sensitivity analyses with and without the exclusion.

**13.** Patient preference is not a clinical exception to eligibility and can be influenced by provider interventions.

**14.** Risk factors that influence outcomes should not be specified as exclusions.

**15.** With large enough sample sizes, small differences that are statistically significant may or may not be practically or clinically meaningful. The substantive question may be, for example, whether a statistically significant difference of one percentage point in the percentage of patients who received smoking cessation counseling (e.g., 74 percent v. 75 percent) is clinically meaningful; or whether a statistically significant difference of \$25 in cost for an episode of care (e.g., \$5,000 v. \$5,025) is practically meaningful. Measures with overall less-than-optimal performance may not demonstrate much variability across providers.

**1. DATA/SAMPLE USED FOR ALL TESTING OF THIS MEASURE**

Often the same data are used for all aspects of measure testing. In an effort to eliminate duplication, the first five questions apply to all measure testing. If there are differences by aspect of testing, (e.g., reliability vs. validity) be sure to indicate the specific differences in question 1.7.

**1.1. What type of data was used for testing?** (Check all the sources of data identified in the measure specifications and data used for testing the measure. Testing must be provided for all the sources of data specified and intended for measure implementation. **If different data sources are used for the numerator and denominator, indicate N [numerator] or D [denominator] after the checkbox.**)

Measure Specified to Use Data From: (must be consistent with data sources entered in S.17)	Measure Tested with Data From:
<input type="checkbox"/> abstracted from paper record	<input type="checkbox"/> abstracted from paper record
<input checked="" type="checkbox"/> claims	<input checked="" type="checkbox"/> claims
<input checked="" type="checkbox"/> registry	<input checked="" type="checkbox"/> registry
<input type="checkbox"/> abstracted from electronic health record	<input type="checkbox"/> abstracted from electronic health record
<input type="checkbox"/> eMeasure (HQMF) implemented in EHRs	<input type="checkbox"/> eMeasure (HQMF) implemented in EHRs
<input type="checkbox"/> other: Click here to describe	<input type="checkbox"/> other: Click here to describe

**1.2. If an existing dataset was used, identify the specific dataset** (the dataset used for testing must be consistent with the measure specifications for target population and healthcare entities being measured; e.g., Medicare Part A claims, Medicaid claims, other commercial insurance, nursing home MDS, home health OASIS, clinical registry).

CROWNWeb and Medicare Claims Data from January 2013 to December 2013

For the Spring 2019 maintenance submission, 2017 CROWNWeb and Medicare claims data were used.

**1.3. What are the dates of the data used in testing?** January 2013 to December 2013

For the Spring 2019 maintenance submission, January – December 2017 data were used.

**1.4. What levels of analysis were tested?** (testing must be provided for all the levels specified and intended for measure implementation, e.g., individual clinician, hospital, health plan)

Measure Specified to Measure Performance of: (must be consistent with levels entered in item S.20)	Measure Tested at Level of:
<input type="checkbox"/> individual clinician	<input type="checkbox"/> individual clinician
<input type="checkbox"/> group/practice	<input type="checkbox"/> group/practice
<input checked="" type="checkbox"/> hospital/facility/agency	<input checked="" type="checkbox"/> hospital/facility/agency
<input type="checkbox"/> health plan	<input type="checkbox"/> health plan
<input type="checkbox"/> other: Click here to describe	<input type="checkbox"/> other: Click here to describe

**1.5. How many and which measured entities were included in the testing and analysis (by level of analysis and data source)?** (identify the number and descriptive characteristics of measured entities)

*included in the analysis (e.g., size, location, type); if a sample was used, describe how entities were selected for inclusion in the sample)*

1,528 facilities with at least 11 PD patients. Public reporting of this measure on DFC or in the ESRD QIP would be restricted to facilities with at least 11 eligible patients for the measure. We have applied this restriction to all the reliability and validity testing reported here.

For the Spring 2019 maintenance submission, 1,984 facilities that had at least 11 eligible patients during January 2017 – December 2017 were included in the analyses. Public reporting of this measure on DFC or in the ESRD QIP would be restricted to facilities with at least 11 eligible patients for the measure to comply with restrictions on reporting of potentially patient identifiable information related to small sample size. We have applied this restriction to all the reliability and validity testing reported here.

**1.6. How many and which patients were included in the testing and analysis (by level of analysis and data source)?** *(identify the number and descriptive characteristics of patients included in the analysis (e.g., age, sex, race, diagnosis); if a sample was used, describe how patients were selected for inclusion in the sample)*

45,554 PD patients in facilities with at least 11 PD patients. 395,589 patient months are included in the analysis.

For the Spring 2019 maintenance submission, 57,969 patients who are from 1,984 facilities with at least 11 eligible patients were included in the analyses. 486,007 patient months are included in the analysis.

**1.7. If there are differences in the data or sample used for different aspects of testing (e.g., reliability, validity, exclusions, risk adjustment), identify how the data or sample are different for each aspect of testing reported below.**

N/A

**1.8 What were the social risk factors that were available and analyzed?** For example, patient-reported data (e.g., income, education, language), proxy variables when social risk data are not collected from each patient (e.g. census tract), or patient community characteristics (e.g. percent vacant housing, crime rate) which do not have to be a proxy for patient-level data.

N/A

---

## **2a2. RELIABILITY TESTING**

**Note:** *If accuracy/correctness (validity) of data elements was empirically tested, separate reliability testing of data elements is not required – in 2a2.1 check critical data elements; in 2a2.2 enter “see section 2b2 for validity testing of data elements”; and skip 2a2.3 and 2a2.4.*

**2a2.1. What level of reliability testing was conducted?** *(may be one or both levels)*

**Critical data elements used in the measure** *(e.g., inter-abstractor reliability; data element reliability)*

*must address ALL critical data elements)*

**Performance measure score** (e.g., *signal-to-noise analysis*)

**2a2.2. For each level checked above, describe the method of reliability testing and what it tests** (*describe the steps—do not just name a method; what type of error does it test; what statistical analysis was used*)

We used January 2013 – December 2013 Claims data to calculate the inter-unit reliability (IUR) for the overall 12 months to assess the reliability of this measure. The NQF-recommended approach for determining measure reliability is a one-way analysis of variance (ANOVA), in which the between and within facility variation in the measure is determined. The inter-unit reliability (IUR) measures the proportion of the measure variability that is attributable to the between-facility variance. The yearly based IUR was estimated using a bootstrap approach, which uses a resampling scheme to estimate the within facility variation that cannot be directly estimated by ANOVA. We note that the method for calculating the IUR was developed for measures that are approximately normally distributed across facilities. Since this measure is not normally distributed, the IUR value should be interpreted with some caution.

*For the Spring 2019 maintenance submission, we followed the same methodology as described above, using January 2017 – December 2017 CROWNWeb and Medicare Claims data.*

**2a2.3. For each level of testing checked above, what were the statistical results from reliability testing?** (*e.g., percent agreement and kappa for the critical data elements; distribution of reliability statistics from a signal-to-noise analysis*)

For reliability we calculated the monthly and annual IUR across the 12 reporting months. As explained above, the method for calculating the IUR was developed for measures that are approximately normally distributed across facilities. IUR=0.910, which is high and suggests 91% of variation in the measure is attributed to between facility variation. The confidence interval is (0.904, 0.917).

*For the Spring 2019 maintenance submission, the annual IUR was 0.858 across 12 reporting months, which is high and suggests 86% of variation in the measure is attributed to between facility variation and approximately 14% to within facility variation.*

**2a2.4 What is your interpretation of the results in terms of demonstrating reliability?** (*i.e., what do the results mean and what are the norms for the test conducted?*)

The IUR suggests this measure is reliable. However, since the distribution of performance scores is skewed, the IUR value should be interpreted with some caution.

*For the Spring 2019 maintenance submission, the IUR again suggests this measure is reliable. However, since the distribution of performance scores is skewed, the IUR value should be interpreted with some caution.*

## 2b1. VALIDITY TESTING

### 2b1.1. What level of validity testing was conducted? (may be one or both levels)

Critical data elements (data element validity must address ALL critical data elements)

Performance measure score

Empirical validity testing

Systematic assessment of face validity of performance measure score as an indicator of quality or resource use (i.e., is an accurate reflection of performance on quality or resource use and can distinguish good from poor performance) **NOTE:** Empirical validity testing is expected at time of maintenance review; if not possible, justification is required.

**2b1.2. For each level of testing checked above, describe the method of validity testing and what it tests (describe the steps—do not just name a method; what was tested, e.g., accuracy of data elements compared to authoritative source, relationship to another measure as expected; what statistical analysis was used)**

Validity was assessed by calculating the Spearman correlation between this measure and the 2013 SMR and SHR.

This measure is also being maintained on the basis of face validity. Use of small solute clearance (urea reduction ratio and more recently Kt/V) as a dialysis quality measure was initially developed and approved by Clinical TEPs in 2006, 2010 (for pediatric HD and PD adequacy), and 2013 which all agreed that this quality measure domain will improve is important in the assessment of the quality of care for dialysis patients. Achieving target Kt/V was finalized for the ESRD QIP beginning with PY 2015, and has been reported on Dialysis Facility Compare since January 2013.

For the Spring 2019 maintenance submission, we followed the same methodology as the previous submission, using January 2017 – December 2017 CROWNWeb and Medicare Claims data and the 2017 versions of the SMR and SHR. Correlations were calculated to assess the association this measure with clinical outcome quality measures expected to be markers of quality care. The measures selected are fully developed and NQF endorsed, and represent an important subset of core clinical quality measures for this patient population.

We expected the following correlations with this measure:

- SMR: We anticipated a negative correlation with this measure
- SHR: We anticipated a negative correlation with this measure

Kt/V is a marker of dialysis adequacy; if targets are not being met, one would anticipate this being reflected in worse patient outcomes such as morbidity and mortality.

This measure is also being maintained on the basis of face validity. Use of small solute clearance (urea reduction ratio and more recently Kt/V) as a dialysis quality measure was initially developed and approved by Clinical TEPs in 2006, 2010 (for pediatric HD and PD adequacy), and 2013; the TEPs all agreed that this quality measure domain is important in the assessment of the quality of care for dialysis patients. Achieving target Kt/V was finalized for the ESRD QIP beginning with PY 2015, and has been reported on Dialysis Facility Compare since January 2013.

**2b1.3. What were the statistical results from validity testing? (e.g., correlation; t-test)**

The Spearman correlation between this measure and the 2013 standardized mortality ratio as measured by the NQF endorsed SMR (NQF 0369) for the same facility is -0.008 (p-value=0.7749). The Spearman correlation between this measure and the 2013 standardized hospitalization ratio as measured by the 2013 SHR (NQF 1463) is -0.139 (p-value <0.0001).

For the Spring 2019 maintenance submission, The Spearman correlation between the PD Kt/V measure and SMR is -0.058, and statistically significant (p=0.01). The Spearman correlation between PD Kt/V measure and SHR is -0.116, and statistically significant (p<.0001).

**2b1.4. What is your interpretation of the results in terms of demonstrating validity? (i.e., what do the results mean and what are the norms for the test conducted?)**

The Spearman correlation estimates indicate higher facility level percentages of patients at the facility that achieve the Kt/V target is associated lower standardized hospitalization, respectively, although the magnitude of the association is low. A very weak association between facility level percentages of patients achieving the Kt/V target and lower standardized mortality was observed and in the expected direction, however the correlation coefficient was not statistically significant.

For the Spring 2019 maintenance submission, The Spearman correlation coefficients indicate higher facility level percentages of patients that achieve the Kt/V target is associated with lower standardized mortality and hospitalization. The direction of the association was as expected and the association was statistically significant.

---

**2b2. EXCLUSIONS ANALYSIS**

NA  no exclusions — skip to section 2b3

**2b2.1. Describe the method of testing exclusions and what it tests (describe the steps—do not just name a method; what was tested, e.g., whether exclusions affect overall performance scores; what statistical analysis was used)**

N/A

**2b2.2. What were the statistical results from testing exclusions? (include overall number and percentage of individuals excluded, frequency distribution of exclusions across measured entities, and impact on performance measure scores)**

N/A

**2b2.3. What is your interpretation of the results in terms of demonstrating that exclusions are needed to prevent unfair distortion of performance results? (i.e., the value outweighs the burden of increased data collection and analysis. Note: If patient preference is an exclusion, the measure must be specified so that the effect on the performance score is transparent, e.g., scores with and without exclusion)**

N/A

## 2b3. RISK ADJUSTMENT/STRATIFICATION FOR OUTCOME OR RESOURCE USE MEASURES

**If not an intermediate or health outcome, or PRO-PM, or resource use measure, skip to section 2b4.**

**2b3.1. What method of controlling for differences in case mix is used?**

- No risk adjustment or stratification**
- Statistical risk model with** [Click here to enter number of factors](#) **risk factors**
- Stratification by** [Click here to enter number of categories](#) **risk categories**
- Other,** [Click here to enter description](#)

**2b3.1.1 If using a statistical risk model, provide detailed risk model specifications, including the risk model method, risk factors, coefficients, equations, codes with descriptors, and definitions.**

N/A

**2b3.2. If an outcome or resource use component measure is not risk adjusted or stratified, provide rationale and analyses to demonstrate that controlling for differences in patient characteristics (case mix) is not needed to achieve fair comparisons across measured entities.**

Risk adjustment is not necessary for this measure. Disparities were examined at the facility level in section 1b.4, and no disparities were found.

**2b3.3a. Describe the conceptual/clinical and statistical methods and criteria used to select patient factors (clinical factors or social risk factors) used in the statistical risk model or for stratification by risk (e.g., potential factors identified in the literature and/or expert panel; regression analysis; statistical significance of  $p < 0.10$ ; correlation of  $x$  or higher; patient factors should be present at the start of care) Also discuss any “ordering” of risk factor inclusion; for example, are social risk factors added after all clinical factors?**

N/A

**2b3.3b. How was the conceptual model of how social risk impacts this outcome developed? Please check all that apply:**

- Published literature**
- Internal data analysis**
- Other (please describe)**

N/A

**2b3.4a. What were the statistical results of the analyses used to select risk factors?**

N/A

**2b3.4b. Describe the analyses and interpretation resulting in the decision to select social risk factors (e.g. prevalence of the factor across measured entities, empirical association with the outcome, contribution of unique variation in the outcome, assessment of between-unit effects and within-unit effects.) Also describe the impact of adjusting for social risk (or not) on providers at high or low extremes of risk.**

N/A

**2b3.5. Describe the method of testing/analysis used to develop and validate the adequacy of the statistical model or stratification approach** (*describe the steps—do not just name a method; what statistical analysis was used*)

*Provide the statistical results from testing the approach to controlling for differences in patient characteristics (case mix) below.*

**If stratified, skip to 2b3.9**

N/A

**2b3.6. Statistical Risk Model Discrimination Statistics** (*e.g., c-statistic, R-squared*):

N/A

**2b3.7. Statistical Risk Model Calibration Statistics** (*e.g., Hosmer-Lemeshow statistic*):

N/A

**2b3.8. Statistical Risk Model Calibration – Risk decile plots or calibration curves:**

N/A

**2b3.9. Results of Risk Stratification Analysis:**

N/A

**2b3.10. What is your interpretation of the results in terms of demonstrating adequacy of controlling for differences in patient characteristics (case mix)?** (*i.e., what do the results mean and what are the norms for the test conducted*)

N/A

**2b3.11. Optional Additional Testing for Risk Adjustment** (*not required, but would provide additional support of adequacy of risk model, e.g., testing of risk model in another data set; sensitivity analysis for missing data; other methods that were assessed*)

N/A

---

## **2b4. IDENTIFICATION OF STATISTICALLY SIGNIFICANT & MEANINGFUL DIFFERENCES IN PERFORMANCE**

**2b4.1. Describe the method for determining if statistically significant and clinically/practically meaningful differences in performance measure scores among the measured entities can be identified** (*describe the steps—do not just name a method; what statistical analysis was used? Do not just repeat the information provided related to performance gap in 1b*)

Differences in measure performance were evaluated separately for each facility using patient level analyses. The proportion of patients with monthly measurement of Adult PD Kt/V, calculated at the year-level, was compared between one facility and the overall national distribution, and repeated for each individual facility.

Note that the monthly based measure is a simple average of binary outcomes across individuals in the facility, for which the binary outcome equals to 0 (failure= not meeting the threshold) if the value is less than 1.7. The differences in proportions can be compared using Fisher's Exact tests or its normal approximation. The yearly based measure, however, is not a simple average of binary outcomes and we instead used a re-sampling based exact test, with re-sampling generated from the population distribution of the patient level outcomes. Due to the non-symmetric structure of the measure

distributions, a one-sided test with significance level 0.025 is used (corresponding to cutoff=0.05 in a two-sided test). To calculate the p-value, we assess the probability that the facility would experience a number of events (i.e., achievement of the Kt/V threshold) more extreme than that observed if the null hypothesis were true, the null hypothesis being that facility level proportions of patient months which achieve the threshold for Adult PD Kt/V will follow the overall national distribution.

For the Spring 2019 submission, we reproduced the significance analysis using data from January – December 2017. We have revised the description of the analysis to be clearer:

Testing was performed on the yearly based performance score. We used a re-sampling based exact test, with re-sampling generated from the population distribution of the patient level outcomes. Note that a one-sided test with significance level 0.025 is used (corresponding to cutoff=0.05 in a two-sided test) due to non-symmetric structure of the measure's distribution. To calculate the p-value, we compute the probability that the facility would experience a number of events (i.e., percentage with hypercalcemia) more extreme than that observed if the null hypothesis were true, with the null hypothesis being that the facility's distribution of hypercalcemia will follow the overall national distribution.

**2b4.2. What were the statistical results from testing the ability to identify statistically significant and/or clinically/practically meaningful differences in performance measure scores across measured entities?** (e.g., number and percentage of entities with scores that were statistically significantly different from mean or some benchmark, different from expected; how was meaningful difference defined)

Proportion of facilities with significant p-values (0-as expected; 1-worse than expected; significance level  $\leq 0.025$ ) is shown as follows:

<u>Category</u>	<u>Number of facilities</u>	<u>Percent of facilities</u>
0	1272	83.25%
1	256	16.75%

For the Spring 2019 maintenance submission, the proportion of facilities with significant p-values (0-as expected; 1-worse than expected; significance level  $\leq 0.025$ ) is shown as follows:

<u>Category</u>	<u>Number of facilities</u>	<u>Percent of facilities</u>
0	1,763	88.86%
1	221	11.14%

**2b4.3. What is your interpretation of the results in terms of demonstrating the ability to identify statistically significant and/or clinically/practically meaningful differences in performance across measured entities?** (i.e., what do the results mean in terms of statistical and meaningful differences?)

Using patient month values for achieving versus not achieving the threshold for Kt/V in adult PD patients, calculated at the year-level, as the performance measure, 1272 (83.25%) facilities have

achieved expected performance, and 256 facilities (16.75%) have performed worse than expected when compared to the overall national proportion of facilities.

In general, higher Kt/V values represent better quality of care. This analysis demonstrates both practical and statistically significant differences in performance across facilities based on their proportion of patient months with a Kt/V that meets the target.

For the Spring 2019 Maintenance submission, 1,763 (88.86%) facilities have achieved expected performance, and 221 facilities (11.14%) have performed worse than expected when compared to the overall national proportion of facilities.

In general, higher Kt/V values represent better quality of care. This analysis demonstrates both practical and statistically significant differences in performance across facilities based on their proportion of patient months with a Kt/V that meets the target.

---

**2b5. COMPARABILITY OF PERFORMANCE SCORES WHEN MORE THAN ONE SET OF SPECIFICATIONS**  
***If only one set of specifications, this section can be skipped.***

**Note:** *This item is directed to measures that are risk-adjusted (with or without social risk factors) OR to measures with more than one set of specifications/instructions (e.g., one set of specifications for how to identify and compute the measure from medical record abstraction and a different set of specifications for claims or eMeasures). It does not apply to measures that use more than one source of data in one set of specifications/instructions (e.g., claims data to identify the denominator and medical record abstraction for the numerator). Comparability is not required when comparing performance scores with and without social risk factors in the risk adjustment model. However, if comparability is not demonstrated for measures with more than one set of specifications/instructions, the different specifications (e.g., for medical records vs. claims) should be submitted as separate measures.*

**2b5.1. Describe the method of testing conducted to compare performance scores for the same entities across the different data sources/specifications (describe the steps—do not just name a method; what statistical analysis was used)**

N/A

**2b5.2. What were the statistical results from testing comparability of performance scores for the same entities when using different data sources/specifications? (e.g., correlation, rank order)**

N/A

**2b5.3. What is your interpretation of the results in terms of the differences in performance measure scores for the same entities across the different data sources/specifications? (i.e., what do the results mean and what are the norms for the test conducted)**

N/A

---

## 2b6. MISSING DATA ANALYSIS AND MINIMIZING BIAS

**2b6.1. Describe the method of testing conducted to identify the extent and distribution of missing data (or nonresponse) and demonstrate that performance results are not biased** due to systematic missing data (or differences between responders and nonresponders) and how the specified handling of missing data minimizes bias (*describe the steps—do not just name a method; what statistical analysis was used*)

Reporting months with missing values are not excluded from this measure. Missing months are used to help define the measure numerator (missing is counted as not meeting the minimum threshold), so introduction of bias from exclusion of missing values is not a consideration for the measure as specified.

**2b6.2. What is the overall frequency of missing data, the distribution of missing data across providers, and the results from testing related to missing data?** (*e.g., results of sensitivity analysis of the effect of various rules for missing data/nonresponse; if no empirical sensitivity analysis, identify the approaches for handling missing data that were considered and pros and cons of each*)

N/A

**2b6.3. What is your interpretation of the results in terms of demonstrating that performance results are not biased** due to systematic missing data (or differences between responders and nonresponders) and how the specified handling of missing data minimizes bias? (*i.e., what do the results mean in terms of supporting the selected approach for missing data and what are the norms for the test conducted; if no empirical analysis, provide rationale for the selected approach for missing data*)

N/A